

ELECTRICAL SHOCK TREATMENT CHART

START THE TREATMENT EVEN THOUGH THE VICTIM APPEARS TO BE DEAD

Method to adopt: When there are no burns on the chest, or abdomen use Scheffer's Prone Pressure method, if there are burns, use Silvester's method.

Silvester's Method: Lay the victim on his back. Unfasten all tight clothing around the chest, waist and stomach. Raise the shoulders of the victim, support him on your knees, while you place your hands, one on each shoulder, and pad the palms of your hands with your fingers. Press down on the shoulders bilaterally, and then pull down on the shoulders bilaterally. Lower the victim so that the padding is in line with his shoulders, and his head falls slightly back. Turn the head on one side, open the mouth and remove tobacco, pen, false teeth, etc. and "One, Two, Three, Four! Make sure that the tongue remains out while treatment is given." Turn the head on the other side, open the mouth and remove tobacco, pen, false teeth, etc. Kneel about 9 inches behind the victim's head as shown in Fig. 4 and raise his arms above his head.

(B) When there are burns on the chest: Grasp the wrists of the victim and press his crossed arms on his abdomen as shown in Fig. 6 and say "One, Two, Three, Four" Next stretch the arms outwards and upwards as described in "A".

- (1) The full cycle of expiration and aspiration should be repeated from 12 to 15 times a minute. On no account must the rhythm be broken.
- (2) Continue artificial respiration until the victim resumes natural breathing, or the doctor pronounces the victim dead.
- (3) When the victim begins to revive, regulate your timing according to his natural breathing.
- (4) Promote warmth by wrapping the victim in a blanket or by the use of hot water bottles.
- (5) Do not allow the victim to get up until he is able to sit up without any strain on the heart.
- (6) When the victim refuses do not allow him to sit up. Encourage the patient with soothing and consoling words. Treat the face.
- (7) Do not allow the victim to drink or attempt to give him any stimulant until he is fully conscious.
- (8) The victim may be removed to hospital only after he has recovered from the immediate effect of the attack.
- (9) Watch the patient carefully, if breathing fails, start artificial respiration again immediately.
- (10) Do not exert more pressure than is necessary, as internal organs may be injured.
- (11) Do not let the pressure increase to the stage of a severe headache.

Note: The above treatment can also be applied to victims of gas drowning after the following preliminary treatment.

[illegible]

है। इस क्रियाओं द्वारा बहुत नियंत्रण दे भी क्रिया करती है। वायव संकुचों होने से अपना बचन जानती है उदा. मेरा, उदा. मेरा बचन बोलना चाहती है। "वायव" को, "वायव" नियंत्रण, ३ में दिखाने में अनुसार इस क्रिया को द्वारा देने की क्रिया करती है। वायव देने से क्रिया देने में उदा. इस बहुत बोलना जानती है। और बचन उदा. मेरे से अच्छी उदा. भीतर उदा. जानती है।

[illegible]

साधारण सुप्रसारी, ज्वार मरुति घोर अन्य प्रायले और दुसरे उपचार

- [illegible]

